

THE MARYLAND CANNABIS ADMINISTRATION'S TECHNICAL AUTHORITY FOR CANNABIS TESTING

EFFECTIVE JANUARY 2024

The Maryland Cannabis Administration (MCA) has developed this technical authority document to define contaminants and corresponding action limits associated with those contaminants in cannabis. This information is intended for use by the independent testing laboratories registered with the MCA.

Table of Contents

Introduction	3
Sampling	4
Collection Procedure for Laboratory Samples	5
Potency	6
Pesticides	7
Residual Solvents	8
Microbiological Impurities	9
Heavy Metals	11
Excipients	13
Stability Testing	13
Appendix A - Cannabis Testing Requirements	14
Appendix B - Definitions	17
Appendix C - Stability Testing Protocol- MCA Licensed Grower	20
Appendix D - Stability Testing Protocol-MCA Licensed Processor	21
Appendix E - Stability Testing Protocol-Edibles	23
Appendix F - Microbiological Quality Control	24
Appendix G- Pathogen Detection Storage Requirements	28
Appendix H- Green Waste Disposal Procedure for Independent Testing Laboratories	29
Appendix I-Standard Method Performance Requirements and Sample Analysis Methods	30
References	33

INTRODUCTION

Analytical testing of cannabis for safety and potency is increasingly recognized as a critical and necessary component of the industry for several reasons (Freeman et al. 2016):

- Laboratory testing minimizes the risk of pesticides, microbes, heavy metals, toxins, and residual solvents from being consumed by an immunocompromised population;
- Quantification of cannabinoid profiles and potency becomes available for the consumer and aids in determining appropriate dosing for individual use; and
- Laboratory testing provides a sense of public safety and product quality for the cannabis tested.

The Maryland Cannabis Administration (MCA), with the assistance of a scientific work group, has established this technical authority to serve as a reference guide for the independent testing laboratories (ITL) to follow when analyzing cannabis. This technical authority has the force and effect of law and must be followed by ITLs pursuant to the Code of Maryland Regulations (COMAR) 14.17.06.02.

Cannabis purity and potency is to be analyzed based on the most current version of the cannabis inflorescence monograph published by the American Herbal Pharmacopeia (AHP), or a scientifically valid methodology that is equal or superior to that of the AHP monograph. This technical authority provides the lists of contaminants and the acceptable tolerances that the ITL is required to report. The tolerances were established following a review of available literature in the cannabis industry as well as references from the International Conference for Harmonisation (ICH) Guideline Q3C on Impurities and the ICH Guideline Q3D on Elemental Impurities Guidance for Industry.

The four categories of contaminants identified for analysis include:

- Pesticides;
- Residual Solvents;
- Microbiological Impurities; and
- Heavy Metals.

In an effective testing program, standardized sampling procedures are an integral component to quality laboratory testing. The data generated from all analytical methods must be consistently reliable and legally defensible. To achieve this, method precision and accuracy measurements should be performed during the sample testing process. This guidance will provide best practices for sample collection by the ITL.

All sampling and analysis described in this guidance shall be conducted by an ITL registered with the MCA and in good standing and accredited to ISO/IEC 17025 by an International Laboratory Accreditation Cooperation (ILAC) recognized third party.

The MCA is committed to evidence-based decision-making when implementing technical guidance for the registered ITL. As research into cannabis use and safety advances, this technical authority will be revised and updated to reflect the state of science as it pertains to the cannabis industry.

SAMPLING

The objective of a sampling procedure is to ensure the proper collection, clear labeling, proper preservation, careful transportation, and storage of samples by trained personnel for laboratory analyses. Collection of the sample is critical as it must be truly representative of the material being analyzed or the results will not be meaningful. ITLs are required to develop a statistically valid sampling method and collect a representative sample from each batch or lot of final product that is adequate to perform the required testing. The amount of sample required for cannabinoid or contaminant testing may vary due to sample matrix, analytical method, and laboratory-specific procedures.

Cannabis sampling procedures play an important role in identifying and/or confirming the integrity of a sample, as well as the completeness of request and chain of custody forms.

To reliably provide the laboratory with a representative sample, standard sampling methods with descriptive steps must be applied with quality and consistency. All sampling must be consistently performed using accepted methodologies. It is the responsibility of the ITL to define a standard operating procedure that minimizes both imprecision and bias and lists chronological steps that ensure a consistent and repeatable method.

When sampling for compliance, all ITLs are required to follow the sampling protocol listed on page 5 of this document, "Collection Procedure for Laboratory Compliance and Retention Samples." In addition, the following sampling guidelines shall be demonstrated by the laboratory when performing sampling at a licensed grower or licensed processor:

- The use of appropriate sampling equipment to avoid contamination;
- The documentation of observations and procedures used during sample collection;
- The use of an aseptic collection technique is required for antimicrobial testing;
- The importance of personal hygiene and use of person protective equipment; and
- The method used by personnel to consistently obtain samples throughout the batch.

(See Appendix A – Cannabis Testing Requirements for information regarding required testing for each sample matrix).

COLLECTION PROCEDURE FOR LABORATORY COMPLIANCE AND RETENTION SAMPLES

Equipment:

1. PPE-Disposable Gloves/Facemask/Shield;
2. Calibrated **Balance**;
3. **Sterile** Sample Collection Vessel;
4. Isopropyl Alcohol; and
5. Required METRC tags.

Procedure:

- 1) Put on disposable gloves to mitigate the risk for contamination of the sample during the collection process.
- 2) Ensure the work surface and **balance** are clean and decontaminated.
- 3) Label an aseptic collection vessel with the appropriate METRC tag and confirm the batch or lot mass.
Do not sample if pertinent information is not available.
- 4) Retrieve the container you will be collecting the sample from and wipe off the lid of the container if applicable.
- 5) For usable cannabis: The minimum sample volume to be collected from each batch is 0.5% of the batch mass. The minimum number of sample increments listed below must be collected for the gross sample (this includes both compliance and retain sample). Withdraw samples from the upper, middle, and lower sections of each container, with the upper section sample being taken from a depth of not less than 10 centimeters. In circumstances where there are 1-10 containers in a batch, collect a sample from all containers. Record the time the sample was collected, any inconsistencies with the sampling plan, and any other remarks that may be relevant to data analysis or quality assurance.

Max Batch Mass	Minimum Compliance Sample Size	Retention Sample Size
<10lbs	10 sample increments totaling 0.5% batch mass	23g
10-20lbs	12 sample increments totaling 0.5% batch mass	23g

For processed products: Each sample must be taken in final product form from randomly chosen positions in the lot. The sample volume collected must meet or exceed minimum volume requirements for all compliance testing performed.

- 6) Place the sample in the appropriate collection vessel, seal, and place to the side.
- 7) Wipe down the balance and work surface using isopropyl alcohol.
- 8) Dispose of gloves.
- 9) Document the appropriate chain of custody information (i.e. sample volume) to be recorded in METRC.

*The following sample collection procedure is based U.S. Pharmacopeia Convention Chemical Tests / 561 Articles of Botanical Origin. 2014 July

POTENCY

Every batch and/or lot of cannabis cultivated and/or processed for transfer to a licensed dispensary must pass the required compliance testing. Potency is analyzed by quantitating the following compounds:

- Δ 9-Tetrahydrocannabinol (Δ 9-THC);
- Δ 8-Tetrahydrocannabinol (Δ 8-THC);
- Total THC (raw plant material only)
- Tetrahydrocannabinolic Acid (THCA);
- Cannabidiol (CBD);
- Cannabidiolic Acid (CBDA);
- The terpenes described in the most current version of the cannabis inflorescence monograph published by the American Herbal Pharmacopeia (AHP);
- Cannabigerol (CBG); and
- Cannabinol (CBN)

To minimize the variability that exists with potency testing of cannabis flower, all method validations and verifications shall use the standard method performance requirements (SMPRs) listed in Appendix I as guidance for acceptability criteria, when applicable. For matrices not listed, the method performance requirements must be as close to the published SMPRs as possible. Additionally, to maintain consistency, the MCA requires that ITLs use the sample preparation and the sample analysis methods listed in Appendix I. MCA accepts a potency variance of +/- 10% due to the heterogeneity of the cannabis plant. Total THC should be reported for raw plant material based on dry weight. The total THC calculation is as follows: $\text{THC} + (0.877 \times \text{THCA})$. This guidance is from the Agriculture Improvement Act of 2018.

*Note: Test samples for potency will consist of a random selection of buds/flower from the analytical sample of cannabis flower collected from a licensee. The laboratory is to maintain procedures for homogenization which are supported through method validation and/or verification. Elevated potency levels will routinely be monitored and confirmed by the MCA. Enforcement action will be taken for laboratories falsely reporting elevated potency levels in METRC and on Certificates of Analysis.

PESTICIDES

Pesticide applicators and applications shall follow State and federal pesticide requirements for any crop protection agent applied. The Maryland Department of Agriculture (MDA) approves crop protection agents available for use on cannabis. For more information on MDA approved crop protection agents, visit the MCA [website](#). MCA's current list of pesticide targets and action limits are documented in Table 1. To minimize the variability that exists with testing of cannabis flower, all method validations shall meet the standard method performance requirements (SMPRs) listed in Appendix I. Cannabis samples with pesticide active ingredients detected above the action level listed below fail, and the product must be destroyed.

Table 1: List of Target Pesticides and Plant Growth Regulators in Parts Per Million (PPM)

Pesticide/PGR	USE	Action Limit (PPM)
Acetamiprid	Insecticide	0.2
Abamectin	Insecticide	0.5
Aldicarb	Insecticide	0.4
Ancymidol	PGR	0.2
Azoxystrobin	Fungicide	0.2
Bifenazate	Insecticide	0.2
Bifenthrin	Fungicide	0.2
Boscalid	Fungicide	0.4
Carbaryl	PGR	0.2
Carbofuran	Insecticide	0.2
Chlorantraniliprole	Insecticide	0.2
Chlorpyrifos	Insecticide	0.2
Clofentezine	Acaricide	0.2
Cyfluthrin	Insecticide	1.0
Daminozide (Alar)	PGR	1.0
DDVP (Dichlorvos)	Insecticide	0.1
Diazinon	Insecticide	0.2
Dimethoate	Insecticide	0.2
Ethephon	PGR	1.0
Etoxazole	Acaricide	0.2
Fenpyroximate	Insecticide	0.5
Fipronil	Insecticide	0.4
Flonicamid	Insecticide	1.0
Fludioxonil	Fungicide	0.4

Pesticide/PGR	USE	Action Limit (PPM)
Flurprimidol	PGR	0.2
Hexythiazox	Ovicide	1.0
Imazalil	Fungicide	0.2
Imidacloprid	Insecticide	0.4
Kresoxim-methyl	Fungicide	0.4
Malathion	Insecticide	0.2
Metalaxyl	Fungicide	0.2
Methiocarb	Insecticide	0.2
Methomyl	Insecticide	0.4
Myclobutanil	Fungicide	0.2
Naled	Insecticide	0.5
Oxamyl	Insecticide	1.0
Pacllobutrazol	PGR	0.4
Permethrin	Insecticide	0.5
Phosmet	Insecticide	0.2
Piperonyl butoxide	Insecticide	1.0
Propiconazole	Fungicide	0.4
Pyrethrins	Insecticide	1.0
Spinosad	Insecticide	0.2
Spiromesifen	Insecticide	0.2
Spirotetramat	Insecticide	0.2
Thiacloprid	Insecticide	0.2
Thiamethoxam	Insecticide	0.2
Trifloxystrobin	Fungicide	0.2

RESIDUAL SOLVENTS

Some producers of cannabis products use solvents to extract and/or concentrate the active ingredients from cannabis. The MCA has adopted a list of target residual solvents based on common extraction and concentration techniques in the industry. Concentration limits are based on the "International Conference for Harmonisation (ICH) Guideline Q3C (R5) on Impurities: Guidelines for residual solvents." The concentration limits listed in ICH Q3C are based on the toxicity of the individual solvent and on the magnitude of exposure to occur from consuming 10 grams of the pharmaceutical. To minimize variability that exists with testing of cannabis flower, all method validations shall include the standard method performance requirements (SMPRs) listed in Appendix I.

Note: No health-based solvent residual limits have been established specifically for cannabis extract or concentrate products. We are uncertain whether the selected action levels for solvents in cannabis products sufficiently protect persons who smoke cannabis. However, the ICH Q3C does assume 100% absorption by any exposure route.

Table 2: Concentration Limits for Residual Solvents in Parts Per Million (PPM)

Solvent	PPM
Heptanes	<5000
Hexanes	<290
Butanes	<5000
Benzene	<2
Toluene	<890
Total Xylenes	<2170
Propanes	<5000
Ethanol	<5000

MICROBIOLOGICAL IMPURITIES

The presence of microbes is common in natural products. It is important to distinguish between organisms ubiquitous in nature and those that are known pathogens. "Indicator tests" don't directly test for pathogens, but instead serve as quality tests or indications that follow-up pathogen testing should be performed (Holmes et al. 2015). Additionally, while microbial and fungal limits are not typically reported as "pass/fail," the MCA has established acceptable limits of detection based on the literature available. The criteria for acceptability in Table 3a and Table 3b (below) list the microbiological impurities and the associated detection limits.

Total Aerobic Microbial Count (TAMC), Total Yeast and Mold Count (TYMC) and Coliform Testing

A registered independent laboratory shall use:

1. An approved AOAC, FDA, or USP validated culture-based method; or
2. Another method approved by MCA.

Pathogen Testing

A registered independent laboratory shall use:

1. An approved AOAC, FDA, or USP validated agar plating method; or
2. (i) Another approved AOAC, FDA, or USP validated method and (ii) agar plating of pathogens.

The laboratory's selected method will require quality controls (positive and negative) performed with each sample set-up, as well as additional criteria identified by each method (e.g., peel plate requires an automatic reader and time stamp). Standard method performance requirements and testing methods available are listed in Appendix I and must be followed by the ITL. See Appendix F - Microbiological Quality Control for additional quality control information and templates. If a pathogen is detected during compliance testing, the ITL should follow protocol listed in Appendix-G- Presumptive Positive Pathogen Detection.

Table 3a: Microbiological Impurities and Accepted Detection Limits in Colony Forming Units (CFU/g) and Parts per Billion (PPB) for flower and processed products.

<u>Microbiological Impurity</u>	<u>CFU/g</u>
Total Aerobic Microbial Count (TAMC)	<100,000
Total Yeast and Mold Count (TYMC)	<100,000
E. coli	<1
Salmonella spp.	"Not Detected"

<u>Mycotoxin</u>	<u>PPB</u>
Aflatoxin B1	<20
Aflatoxin B2	<20
Aflatoxin G1	<20
Aflatoxin G2	<20
Ochratoxin A	<20

Table 3b: Microbiological Impurities and Accepted Detection Limits in Colony Forming Units (CFU/g) for Edibles Products.

<u>Microbiological Impurity</u>	<u>CFU/g</u>
Total Coliforms	<100
Shiga Toxin producing E. coli (STEC)	"Not Detected"
Salmonella, spp	"Not Detected"
L. monocytogenes	"Not Detected"
E. coli*	<1 or "Not Detected"

*E. coli should be recorded as <1 CFU/g for quantitative analysis and "Not Detected" for qualitative analysis.

Water activity (A_w) is a measure of the available water that can be utilized for microbiological growth. A_w ranges from 0 to 1 with microbial growth unlikely below A_w 0.6. Most cannabis is dried and cured to a final water activity level of A_w 0.3-0.6, and most pathogens cannot grow below A_w 0.9 (Holmes et al. 2015). Water activity, or the moisture of the cannabis flower in units, measured below A_w 0.65 will safeguard cannabis products against microbial growth during storage and before sale. To maintain consistency, the MCA requires that ITLs use the sample analysis method listed in Appendix I for water activity.

Table 3c. Acceptable water activity limits for cannabis flower and edible cannabis products. Liquid edible products are excluded from water activity testing.

<u>Water Activity</u>	<u>(A_w)</u>
Flower products	<.65
Edible cannabis products	<.85

HEAVY METALS

Elemental impurities do not provide any therapeutic benefit to the cannabis patient or consumer. Because of their high degree of toxicity, arsenic, cadmium, chromium, lead, and mercury rank among the priority metals that are of public health significance (Tchounwou P et al. 2012). The MCA requires an ITL to test for heavy metal presence in cannabis. Table 4a lists the five heavy metals required in compliance testing and their associated action limits based on a 5 gram/day consumption for inhalation limits and a 10 gram/day consumption for oral limits. Table 4b lists the four heavy metals required in contaminant testing for edible cannabis products and their associated concentration limits based on a 10 gram/day consumption. To minimize variability that exists with testing of cannabis flower, all method validations shall meet the standard method performance requirements (SMPRs) listed in Appendix I. For consistency, the MCA requires that ITLs use the sample analysis method listed in Appendix I for detection and quantification of heavy metals in cannabis flower and processed products.

Note: The permitted daily exposure (PDE) for heavy metals is based on the Q3D Elemental Impurities Guidance for Industry.

Table 4a: Heavy Metals and Associated Concentration Limits in Parts Per Million (PPM) for Flower and Processed Products.

Heavy Metal	PPM (Inhalation)	PPM (Oral)
Lead	<1.0	<0.5
Arsenic	<0.4	<1.5
Mercury	<0.2	<3.0
Cadmium	<0.4	<0.5
Chromium	<0.6	<1070.0

Table 4b: Heavy Metals and Associated Concentration Limits in Parts Per Million (PPM) for Edible Cannabis Products.

Heavy Metal	PPM (Oral)
Lead	<0.5
Arsenic	<1.5
Mercury	<3.0
Cadmium	<0.5

EXCIPIENTS

The presence of any residual solvent or processing chemical shall not exceed the levels provided in this document. On November 15, 2019, the Commission issued [Bulletin 2019-013](#) banning the use of Vitamin E Acetate (VEA) as a processing chemical in the production of cannabis vaping products and requiring VEA screening be performed on all vaping products (see Appendix 1). VEA detection in vape samples that exceed 0.7% by weight will be cause for product destruction.

STABILITY TESTING

Stability testing is to be performed at 6-month intervals. The purpose of stability testing is to provide evidence on how the quality of a drug substance varies with time under the influence of a variety of environmental factors (ICH 2003).

The ITL must have policies and procedures established for the collection of stability and retention samples and the analysis of stability testing samples.

The stability testing required will include:

- Cannabinoid content; and
- Microbiological impurities.

Findings of the stability studies must be reported to the MCA through the METRC tracking system to ensure cannabis purity and potency are maintained throughout the storage process without significant change. Significant change for cannabis is defined as failure to meet the tolerances listed in this technical guidance for purity. Stability studies protocol may change as the industry evolves. Current protocols are listed below.

Stability testing protocol for MCA licensed growers is available in Appendix C – Stability Testing Protocol – MCA Licensed Grower.

Stability testing protocol for MCA licensed processors is available in Appendix D – Stability Testing Protocol – MCA Licensed Processor.

Stability testing protocol for edibles products is available in Appendix E – Stability Testing Protocol - Edibles.

APPENDIX A - Cannabis Compliance Testing Requirements

	Raw Plant Material (Buds, Shake/ Trim, Prerolls)	Concentrate (Solvent/Non -Solvent Based)	Infused Non- Edible	Inhalable/ Vape Concentrate	Infused Edible	Infused Edible (Capsule)	Infused Liquid Edible	Infused Edible (Exempt)	Tincture for Oral Administration	External Hemp (Extract/Raw Plant Material)
Moisture Content	√									
Potency Analysis	√	√	√	√	√	√	√	√	√	√
Terpene Analysis	√	√	√	√					√	
Foreign Matter Inspection	√	√	√	√	√	√	√	√	√	
Microbial Screen (TYMC, TAMC)	√	√	√	√					√	
Mycotoxin Screen	√	√	√	√	√	√	√	√	√	
Water Activity	√				√	√		√		
Heavy Metal Screen	√	√	√	√	√	√	√	√	√	
Residual Solvent Test		√	√	√					√	
Pesticide Residue Analysis	√	√	√	√					√	
Vitamin E Acetate				√						
Shiga Toxin Producing E. Coli					√	√	√	√		
Salmonella, spp.	√	√	√	√	√	√	√	√	√	
Total Coliform					√	√	√	√		
E. coli	√	√	√	√	√	√	√	√	√	
L. monocytogenes					√	√	√	√		

APPENDIX A(1)- PRODUCT CATEGORIES AND COA REPORTING REQUIREMENTS

METRC Product Category	COA Reporting Requirement
Raw Plant Material (Buds, Shake/trim, prerolls)	Cannabinoids/Terpenes-Quantitative Values (%) Moisture Content-Quantitative Value (%) Water Activity-Quantitative Value (A_w) Foreign Matter Inspection-Quantitative Value (%) Pesticides-Pass/Fail Microbial Screen (TYMC, TAMC)-Pass/Fail Salmonella-Pass/Fail E. coli-Pass/Fail Mycotoxin Screen-Pass/Fail Heavy Metals Screen-Pass/Fail
Concentrates	Cannabinoids/Terpenes-Quantitative Values (mg/g) Foreign Matter Inspection-Quantitative Value (%) Microbial Screen (TYMC, TAMC)-Pass/Fail Salmonella-Pass/Fail E. coli-Pass/Fail Mycotoxin Screen-Pass/Fail Heavy Metals Screen-Pass/Fail Residual Solvents-Pass/Fail Pesticides-Pass/Fail
Infused Non-edible.	Cannabinoids/Terpenes-Quantitative Values (mg/g) Foreign Matter Inspection-Quantitative Value (%) Microbial Screen (TYMC, TAMC)-Pass/Fail Salmonella-Pass/Fail E. coli-Pass/Fail Mycotoxin Screen-Pass/Fail Heavy Metals Screen-Pass/Fail Residual Solvents-Pass/Fail Pesticides-Pass/Fail
Vape Cart	Cannabinoids/Terpenes-Quantitative Values (mg/g) Foreign Matter Inspection-Quantitative Value (%) Microbial Screen (TYMC, TAMC)-Pass/Fail Salmonella-Pass/Fail E. coli-Pass/Fail Mycotoxin Screen-Pass/Fail Residual Solvents-Pass/Fail Pesticides-Pass/Fail Vitamin E Acetate-Pass/Fail Heavy Metals Screen-Pass/Fail
Infused Edible	Cannabinoids-Quantitative Values (mg/g) Water Activity-Quantitative Values (A_w) Foreign Matter Inspection-Quantitative Value (%) Total Coliforms-Pass/Fail Shiga toxin producing E. coli-Pass/Fail Salmonella-Pass/Fail E. coli-Pass/Fail L.monocytogenes-Pass/Fail Mycotoxin Screen-Pass/Fail Heavy Metals Screen-Pass/Fail

Infused Edible (Capsule)	Cannabinoids-Quantitative Value (mg/g) Water Activity-Quantitative Value (A_w) Foreign Matter Inspection-Quantitative Value (%) Mycotoxin Screen-Pass/Fail Heavy Metals Screen-Pass/Fail Total Coliforms-Pass/Fail Shiga toxin producing E. coli-Pass/Fail Salmonella-Pass/Fail E. coli-Pass/Fail L.monocytogenes-Pass/Fail
Infused Liquid Edible	Cannabinoids-Quantitative Value (mg/g) Foreign Matter Inspection-Quantitative Value (%) Mycotoxin Screen-Pass/Fail Heavy Metals Screen-Pass/Fail Total Coliforms-Pass/Fail Shiga toxin producing E. coli-Pass/Fail Salmonella-Pass/Fail E. coli-Pass/Fail L.monocytogenes-Pass/Fail
Exempt Edible Product Exempt Liquid Edible Product	Cannabinoids-Quantitative Value (mg/g) Water Activity-Quantitative Value (A_w) (only for non-liquid products) Foreign Matter Inspection-Quantitative Value (%) Mycotoxin Screen-Pass/Fail Heavy Metals Screen-Pass/Fail Total Coliforms-Pass/Fail Shiga toxin producing E. coli-Pass/Fail Salmonella-Pass/Fail E. coli-Pass/Fail L.monocytogenes-Pass/Fail
Tincture for Oral Administration	Cannabinoids/Terpenes-Quantitative Value (mg/g) Foreign Matter-Quantitative Value (%) Mycotoxin Screen-Pass/Fail Heavy Metals Screen-Pass/Fail Microbial Screen (TYMC, TAMC)-Pass/Fail Salmonella-Pass/Fail E. coli-Pass/Fail Pesticides-Pass/Fail Residual Solvents-Pass/Fail

APPENDIX B - DEFINITIONS

Administration- The Maryland Cannabis Administration established under Alcoholic Beverages and Cannabis Article, 36-201, Annotated Code of Maryland.

Batch -

- (a) All of the plants of the same variety of cannabis that have been:
 - (1) Grown, harvested, and processed together; and
 - (2) Exposed to substantially similar conditions throughout cultivation and processing.
- (b) Includes all of the processed materials produced from those plants.

Cannabis-

- (a) The plant cannabis sativa L. and any part of the plant, including all non-synthetically derived, extracts, cannabinoids, isomers, acids, salts, and salts of isomers, whether growing or not, with a delta-9-tetrahydrocannabinol concentration greater than 0.3 percent on a dry weight basis.
- (b) Includes cannabis products.
- (c) Does not include hemp or hemp products, as defined in the Agriculture Article, 14-101, Annotated Code of Maryland.

Cannabis Product-

- (a) A product that is composed of cannabis, cannabis concentrate, cannabis extract, or any other ingredient and is intended for use or consumption.
- (b) Includes any product produced and regulated under subtitle 14.17, including:
 - (1) Cannabis vaporizing devices;
 - (2) Concentrated cannabis products;
 - (3) Edible cannabis products; and
 - (4) Usable cannabis products.
- (c) Does not include a home cultivation product.

Capsule- A solid preparation containing a single serving of tetrahydrocannabinol or other cannabinoid that:

- (a) Is intended to be swallowed whole;
- (b) Not formulated to be chewable, dispersible, effervescent, orally disintegrating, used as a suspension, or consumed in a manner other than swallowed whole; and
- (c) Does not contain any added natural or artificial flavor or sweetener.

Chain of Custody - The chronological documentation showing the collection, custody, control, transfer, analysis, and disposition of a sample.

Concentrated Cannabis Product-

- (a) A product derived from cannabis that has undergone a process to concentrate one or more active cannabinoids.
- (b) Concentrated cannabis products include:
 - (1) Kief;
 - (2) Hashish;
 - (3) Bubble hash;
 - (4) Oil;
 - (5) Wax;
 - (6) Shatter;

- (7) Resin; or
 - (8) Any other product produced by extracting cannabinoids from the plant using solvents, carbon dioxide, heat, screens, presses, or steam distillation.
- (c) Does not include any cannabis vaporizing device as defined in 14.17.

CFU/g - Colony forming units per gram. Refers to a measure of the amount of living bacteria per given amount (1 gram) of a sample.

Edible Cannabis Product-

- (a) Means a cannabis product intended for human consumption by oral ingestion, in whole or in part.
- (b) Includes a cannabis product that dissolves or disintegrates in the mouth.
- (c) Does not include any concentrated cannabis products, infused non-edible cannabis products, or capsules or tinctures that do not contain any food or food ingredients.

Green Waste-unauthorized misbranded, contaminated, unused, surplus, returned, or out of date cannabis or product containing cannabis.

Independent Testing Laboratory - A facility, entity, or site that is:

- (a) Registered with the Administration to perform tests on cannabis or cannabis products;
- (b) Independent of any entity licensed under Alcoholic Beverages and Cannabis Article, 36-401 to grow, process or dispense cannabis; and
- (c) Accredited as operating to International Organization for Standardization (ISO) standard 17025 by and accreditation body:
 - (1) Operating in accordance with ISO standard ISO/IEC 17011; and
 - (2) That is a signatory to the International Laboratory Accreditation Cooperation (ILAC) Mutual Recognition Arrangement (MRA).

Infused Non-edible- means ointment, salve, suppository, dermal patch, cartridge, or any other product containing cannabis that has been processed so that the dried leaves and flowers are integrated into other material that is not intended for human consumption by oral ingestion.

Liquid edible product-

- (a) An edible cannabis product that is a liquid beverage or liquid food-based product for which the intended use is oral consumption.
- (b) Excludes a tincture as defined in 14.17.

Lot - All of a cannabis finished product that is uniform, that is intended to meet specifications, and that is manufactured, packaged, or labeled together during a specified time period according to a single lot record.

METRC –Marijuana Enforcement Tracking Regulation and Compliance system.

Qualitative - Relating to, measuring, or measured by the quality of something rather than its quantity.

Quantitative - Relating to, measuring, or measured by the quantity of something rather than its quality.

Seed-to-sale tracking system-A software system procured by the Administration that tracks cannabis from either the seed or immature plant stage, until the cannabis is sold to a patient, caregiver, or consumer.

Representative Sample - A sample obtained according to a sampling procedure designed to ensure that the different parts of a batch or lot or the different properties of a batch or lot are proportionally represented.

Sample - An amount of cannabis collected by laboratory personnel from a licensee and provided to an independent testing laboratory for testing.

Solvent - A substance that can dissolve another substance, or in which another substance is dissolved, forming a solution.

Target Analyte - A chemical the laboratory must test for to see if it is present in cannabis.

Tetrahydrocannabinol or "THC"-unless otherwise specified means any:

- (a) Tetrahydrocannabinol, including delta-8-tetrahydrocannabinol, delta-9-tetrahydrocannabinol, and delta-10-tetrahydrocannabinol, regardless of how derived;
- (b) Other cannabinoid, other than cannabidiol that the Administration determines to cause intoxication; and
- (c) Other chemically similar compound, substance, derivative, or isomer of tetrahydrocannabinol, as identified by the Administration.

Tincture-a solution that is:

- (a) Dissolved in alcohol, glycerin, or vegetable oil; and
- (b) Distributed in a dropper bottle of four ounces or less.

Usable Cannabis -

- (a) The dried leaves and flowers of the cannabis plant.
- (b) Does not include seedlings, seeds, stems, stalks or roots of the plant or the weight of any noncannabis ingredients combined with cannabis, such as ingredients added to prepare a topical administration.

Usable Cannabis Product-

- (a) A prepackaged product containing usable cannabis.
- (b) Includes:
 - (1) A pre-rolled amount of usable cannabis;
 - (2) Securely stored, sealed, and labeled amount of usable cannabis; and
 - (3) Any other type or amount of usable cannabis that has been wrapped, rolled, or otherwise encased for the purposes of smoking.

Water Activity - The partial vapor pressure of water in a substance divided by the standard state partial vapor pressure of water.

APPENDIX C - STABILITY TESTING PROTOCOL (GROWER)

Stability testing shall be performed for each released batch of usable cannabis. This document outlines the required protocol to be followed by MCA licensed growers and MCA registered ITLs performing the stability studies.

Definitions:

Testing Panel - Each stability sample is to be tested for a) Micro-organisms; and b) Potency to ensure product potency and purity and provide support for expiration dating.

Stability Sample – 12 grams of material stored in routine conditions by the licensed grower to allow for collection of testing samples at all time points.

Testing Sample – 3 grams collected from the stability sample to be collected by, homogenized, and analyzed by the ITL for each time point.

Time Point – The 6-month interval when testing should occur (0, 6, 12 and 18 months).

Homogenization – Manipulation of a product by mixing, and/or grinding, to obtain equal distribution of all components or ingredients with the goal of reducing variability.

Stability Testing Goals:

The design will assess:

- Degradation of cannabinoids in usable cannabis products over an 18-month period when held at routine storage conditions at a licensed cultivation facility.
- Levels of bacterial/fungal growth in usable cannabis products over an 18-month period when held at routine storage conditions at a licensed cultivation facility.

Stability Testing Protocol Requirements:

1. Stability testing shall be performed for each unique strain of cannabis. If material produced is to be distributed/sold as unique products (flower, trim, kief) each of these products shall constitute a batch and must be tested individually as potency, microbiological activity and environmental impact on stability may vary between product forms.
2. The licensed grower shall be responsible for stability sample storage, and selection of the ITL to perform stability testing.
3. The ITL shall be responsible for the collection of the stability and testing samples, analysis, and submission of stability testing data into METRC.
4. Each stability sample shall contain 12 grams of material to allow the ITL to collect a 3-gram testing sample at each of the four time points. Failure to generate sufficient data for analysis may require repeating the missing time point/testing and potentially the full protocol. In cases where insufficient material to complete full testing is available (kief, trim) from a single batch a modified protocol to assess the stability of these products shall be proposed by the licensed grower for approval by the MCA.
5. Stability samples shall be uniquely identified, clearly labeled "For Stability Testing Only" and stored in the same environmental conditions as product intended for sale. Care shall be taken to keep the sample segregated from other products to avoid potential contamination of study samples.
6. The ITL shall collect a testing sample of 3 grams from the stability sample at each time point. In cases where the product is packaged in volumes lower than what is required by the laboratory for testing multiple packages of a product from the same batch may be used to produce a single, homogenized sample for testing. These packages shall be collected by the independent testing laboratory and combined into a single sample at the time of testing.
7. Testing samples are to be collected and analyzed by the ITL at 0, 6, 12 and 18 months.
8. Testing performed at T0 is the full compliance panel. Testing performed at T6, T12, and T18 will consist of potency, TYMC, TAMC, E. coli, and Salmonella.
9. Testing results for all time points shall be generated within 14 calendar days of the date of the time point to be measured.
10. Each testing sample must be homogenized consistent with the laboratory's standard operating procedures.
11. Laboratory methodology shall be consistent throughout the study. Changes to technology or protocols throughout the study require approval from MCA.
12. The ITL shall provide all data electronically to the MCA via an electronic reporting [portal](#) within 30 calendar days of the measured time point.

APPENDIX D - STABILITY TESTING PROTOCOL (PROCESSOR)

Licensed Processor Stability Testing Protocol

Stability testing shall be performed for each released lot of processed cannabis. This document outlines the required protocol to be followed by MCA licensed processors and MCA registered ITLs performing the stability studies.

Definitions:

Cannabis-Infused Product – Oil, wax, ointment, salve, tincture, capsule, suppository, dermal patch, cartridge, or other product containing cannabis concentrate or usable cannabis that has been processed so that the dried leaves and flowers are integrated into other material.

Lot – All of a cannabis finished product that is uniform, that is intended to meet specifications, and that is manufactured, packaged, or labeled together during a specified time period according to a single lot record.

Testing Panel - Each testing sample is to be tested for a) Micro-organisms; and b) Potency.

Stability Sample – Sufficient material stored in routine conditions by the licensed processor to generate testing samples at all time points.

Testing Sample – Sample to be collected from the stability sample by the ITL sufficient to complete the testing panel for each time point.

Time Point – 6-month interval when testing should occur (0, 6, 12 and 18 months).

Homogenization – Manipulation of a product by mixing, to obtain equal distribution of all components or ingredients with the goal of reducing sample variability.

Stability Testing Goals:

The design must assess:

- Degradation of cannabinoids in cannabis processed products over an 18-month period when held at routine storage conditions at a licensed processing facility.
- Levels of bacterial/fungal growth in cannabis processed products over an 18-month period when held at routine storage conditions at a licensed processing facility.

Stability Testing Protocol Requirements:

1. Stability testing shall be performed for each unique cannabis-infused product. Each product with a unique strain, terpene/cannabinoid profile or delivery method shall be tested independently as potency, microbiological activity and environmental impact on stability may vary between product forms.
2. The licensed processor shall be responsible for stability sample storage and selection of the ITL to perform stability testing.
3. The ITL shall be responsible for the collection of the stability and testing samples, analysis, and submission of stability testing data into METRC.
4. Each stability sample shall contain sufficient material to allow the independent testing laboratory to collect a testing sample at each of the four time points sufficient to complete the testing panel. Failure to generate sufficient data for analysis may require repeating the missing time point/testing and potentially the full protocol.
5. Stability samples shall be uniquely identified, clearly labeled "For Stability Testing Only" and stored in the same environmental conditions as product intended for sale. Care shall be taken to keep the sample segregated from other products to avoid potential contamination of study samples.
6. The ITL shall collect a testing sample from the stability sample at each time point sufficient to complete the full testing panel. In cases where the product is packaged in volumes lower than what is required by the laboratory for testing multiple packages of a product from the same batch may be used to produce a single, homogenized sample for testing. These packages shall be collected by the ITL and combined into a single sample at the time of testing.

7. Testing samples are to be collected and analyzed by the independent testing laboratory at 0, 6, 12 and 18 months. Testing performed at T0 is the full compliance panel. Testing performed at T6, T12, and T18 will consist of potency, TYMC, TAMC, E. coli, and Salmonella.
8. Testing results for all time points shall be generated within 14 calendar days of the date of the time-point to be measured.
9. Laboratory methodology shall be consistent throughout the study. Changes to technology or protocols throughout the study require approval from MCA.
10. When possible, each sample is to be homogenized at the time of testing by the ITL consistent with the laboratory's standard operating procedure.
11. ITLs shall provide all data [electronically](#) to the MCA within 30 calendar days of the measured time point.

APPENDIX E - STABILITY TESTING PROTOCOL (EDIBLES)

Edible Products Shelf Stability Study

Shelf-life testing shall be performed for each unique edible cannabis product available for patient and adult-use consumption. This document outlines the required protocol to be followed by MCA licensed processors and the MCA registered ITLs performing testing. The protocol consists of 10 individual product samples being analyzed for content uniformity as well as a 12-week time monitoring product potency, water activity, and microbiological contaminants.

Content Uniformity Requirements (Time point 0):

1. The licensed processor shall randomly select 10 individual samples of unique edible cannabis products in final form from available production lots, ensuring all production lots available have been represented. These samples must be transferred to an ITL for required testing. Compliance testing performed at T0 will satisfy baseline water activity and microbiological data points. The ITL is responsible for randomly sampling for compliance.
2. The ITL shall visually inspect each sample for foreign matter, odor, and general appearance.
3. Following visual inspection, the samples must each be tested for cannabinoid content. Acceptable content uniformity shall fall within +/- 10%.
4. Following completion of testing, results shall be uploaded directly into METRC by the ITL. Additionally, laboratories should submit testing data [electronically](#) to MCA.

Stability Requirements (Time points 1-3):

Following the initial content uniformity testing there will be three additional time points to test: T(1) at 4 weeks, T(2) at 8 weeks, and T(3) at 12 weeks.

1. The licensed processor should randomly select 3 samples (beginning, middle, and end) from each unique production lot at stated time points.
2. The ITL shall visually inspect each sample for foreign matter, odor, and general appearance. Following the visual inspection, the samples must be homogenized and tested for the following:
 - Microorganisms;
 - Water activity; and
 - Cannabinoid content.
3. Testing results must be uploaded directly into METRC by the ITL. Additionally, laboratories shall submit testing data [electronically](#) to MCA.

APPENDIX F - MICROBIOLOGICAL QUALITY CONTROL

Quantitative quality controls are required to quantitate microbes. ITLs shall run quality controls (QC) each time samples are set up. QC must mimic the sample analysis and needs to run through every incubation period during every run (i.e. a broth base analysis must include a broth-based QC, and a plate-based analysis must include a plate-based QC).

Quality Control (QC) Templates are available on cannabis.maryland.gov.

F(1). Quantitative Analysis Control Chart - Broth-based QC

+Control=E.coli, -Control=S. aureus, Sterility Control=Media blank

Test Controls		E. coli ATCC 25922	E. aerogenes ATCC 13048	S. aureus ATCC 25923	Sterility Control	Initial/ Date
LST Control Results		XX	XX	XX	XX	
		X	X	X	X	
EC Control Results						
BGB Control Results						

Temp Incubated _____ °C Time/Date _____ Initials _____

Quantitative QC Petri film/charm controls

Test Controls	E. coli ATCC 25922	S. aureus ATCC 25923	Sterility Control	Initial/ Date
charm/petri film plates	pos control count	neg control		

Temp Incubated _____ °C Time/Date _____ Initials _____

Aerobic Bacteria Count

Aerobic bacteria plate counts controls

<u>PCA Control Plate</u>	<u>Colony Count</u>	<u>Initial/Date</u>
15 min Air Exposure Plate		
Glass Ware		
PCA		
Butterfield's phosphate-buffered/buffer used		
Positive Quantitative QC value		

Temp Incubated _____ °C Time/Date _____ Initials _____

Certified Reference Material/Reference Material Used During Analysis

CRM	Lot #	ATCC or NCTC #	Generation	Expiration Date
Escherichia coli				
Enterobacter aerogenes				
Staphylococcus aureus				
Proteus mirabilis				

F(2). Qualitative Quality Control

Quality Control (QC) performed for qualitative analysis must include a Sterility Control, Negative Control, and a Positive Control with each RUN or at a MINIMUM every time you set up samples for that day. The QC must simulate the samples during each phase. If the sample tested is going through an incubation at a specific temperature, then the QC must mirror it on the same medium. Please see the chart below which shows Salmonella as a positive control, E coli as a negative control and Media blank as a sterility control.

Qualitative Analysis Control Chart

+Control=Salmonella, -Control=E.coli, Sterility Control=Media blank

Test Controls	Salmonella sp.	E. coli	Sterility Control	Initial/ Date
RV Broth				
Tetrathionate Broth				
XLD Agar				
Hektoen Agar				
Wilson Blair Agar				
TSI /LIA/BAP				

Initials/Date: _____ Incubator temperature _____ Water bath temperature _____

Certified Reference Material/Reference Material

CRM	ATCC or NCTC#	Lot#	Generation	Expiration Date
Salmonella species				
Escherichia coli				

APPENDIX G-PRESUMPTIVE POSITIVE PATHOGEN DETECTION

If an ITL identifies a pathogen (E. coli, Salmonella, or Listeria) during routine compliance testing, the following steps should be taken within 24 hours of the presumptive positive:

1. Perform the confirmation steps listed in the chart below for each organism detected.
2. Notify the MCA via phone and email (mca.labs@maryland) of the confirmed pathogen positive.
3. Enter all failed test results into METRC.
4. Notify the licensee and coordinate sample pick with the MCA.
 - a. Refrigerate selective agar plates, the original enrichment broth, and blood plates at 2-8° Celsius until pickup by MCA.
5. Perform environmental swab testing of the licensee's facility after a presumptive positive.
6. Perform environmental swab testing of the licensee's facility after it has been decontaminated.
7. Notify the MCA with the environmental swab testing results.

PATHOGEN DETECTED (AGAR PLATING ONLY)	CONFIRMATION STEPS REQUIRED
E. COLI	API 20E
STEC	API 20E; IF CONFIRMED AS E. COLI LATEX AGGLUTINATION FOR 0157 OR NON-0157, AND/OR QPCR
L.MONOCYTOGENES	API LISTERIA, AND/OR LATEX AGGLUTINATION
SALMONELLA SPP.	API 20E, AND/OR QPCR MAY BE PERFORMED

APPENDIX H- GREEN WASTE DISPOSAL PROCEDURE FOR INDEPENDENT TESTING LABORATORIES

This standard operating procedure provides a standardized method of disposal for cannabis green waste at MCA registered independent testing laboratories. The procedure ensures accountability for cannabis green waste by establishing appropriate documentation and destruction processes. MCA requires registered independent testing laboratories to implement the Green Waste Disposal Procedure upon adoption of Revision 5.0 of this technical authority.

Procedure:

1. Following conclusion of the lab's identified retention period, all waste shall be documented on the Cannabis Green Waste Log attached to this Standard Operating Procedure. The log shall be available for immediate review upon request by MCA personnel. The log must include the following information:
 - A. Date and time the waste was entered into a waste container;
 - B. Product name;
 - C. Last 9 digits of metric tag number;
 - D. Product weight to be green wasted measured in applicable units (i.e. grams, each);
 - E. Agent entering into Waste Log;
 - F. Date and time of disposal and removal from the facility and into a commercial waste bin for pickup;
 - G. Method of disposal (Ex: kitty litter, mulch, bleach);.
 - H. Agent or manager disposing of waste; and
 - I. Agent verifying disposal of waste.
2. All waste shall immediately be rendered unusable, entered onto the waste log, and placed into the waste container. This action must be clearly captured on video.
 - A. Flower/dry leaf waste shall be ground to the smallest possible degree and mixed with a non-cannabis product in a 50:50 ratio (minimum).
(Examples of non-cannabis products include alcohol, bleach, any other solvent that renders it non-useable, kitty litter, mulch, dirt, or other loose non-consumable material that will render the cannabis non-useable).
 - B. Non-flower/non-dry leaf waste shall be emptied into a non-consumable product for disposal.
3. Final destruction shall occur no later than 7 days after the waste is entered onto the Cannabis Green Waste Log and placed in a designated commercial waste bin for pick up and physical removal from the lab's inventory. All waste being disposed must be captured on video and will require verification from two laboratory agents documented on the Cannabis Green Waste Log.
4. All entries in the Cannabis Green Waste Log shall be printed legibly and be consistent with METRC green waste entries. To download a printable copy of the Cannabis Green Waste Log, please click [here](#).

APPENDIX I- METHOD PERFORMANCE REQUIREMENTS AND PERFORMANCE TESTED METHODS AVAILABLE FOR USE

*SMPR's and PTM's will be revised annually. PTM's published in the interim must be approved for use by the MCA. Method validations are required when an independent testing laboratory is developing its own method for use. The following criteria must be provided and approved by MCA in advance of the method being utilized as well as demonstrated by the ITL:

- Sensitivity
- Selectivity
- Repeatability
- Reproducibility
- Robustness
- Accuracy
- Linearity
- LOD
- LOQ

Method verifications are required to verify that an independent testing laboratory can meet the quality control requirements of a reference or validated method. Quality control requirements of the method include but are not limited to the following:

- Limit of detection and quantitation studies
- Initial and continuing calibration verification, as defined by the method
- Control spikes and/or fortified blanks
- Passing proficiency testing samples in the appropriate matrix (if commercially available)

POTENCY:

For method validations, incorporate the Standard Method Performance Requirements (SMPRs) listed below:

- **Dried Plant Material: AOAC SMPR 2017.002**
- **Concentrates: AOAC SMPR 2017.001**

- **Edible Chocolate: AOAC SMPR 2017.019**

Sample Analysis:

- **Quantitation of Cannabinoids in Cannabis Dried Plant Materials, Concentrates, and Oils AOAC 2018.11**

PESTICIDES:

For method validations, incorporate the Standard Method Performance Requirements (SMPRs) listed below:

- **Identification and Quantification of Selected Pesticide Residue in Dried Cannabis Flower: AOAC SMPR 2018.011**

RESIDUAL SOLVENTS:

For method validations, incorporate the Standard Method Performance Requirements (SMPRs) listed below:

- **Identification and Quantitation of Selected Residual Solvents in Cannabis-Derived Materials: AOAC 2019.002**

MICROBIOLOGICAL IMPURITIES:

For method validations, incorporate Standard Method Performance Requirements (SMPRs) listed below:

- **Detection of Salmonella species in Cannabis and Cannabis Products: AOAC 2020.002**
- **Detection of Shiga Toxin-Producing Escherichia coli in Cannabis and Cannabis Products: AOAC 2020.012**
- **Viable Yeast and Mold Count Enumeration in Cannabis and Cannabis Products: AOAC 2021:009**
- **Mycotoxin Screening Technique in Cannabis Plant Material and Cannabis Derivatives: AOAC 2020.013**

Sample Analysis:

- **Yeast and Mold Counts in Foods and Dried Cannabis Flower: AOAC 997.02**
- **3M Petrifilm Rapid Yeast and Mold Plate Count**
- **TEMPO Yeast & Mold**
- **TEMPO AC (Aerobic Count)**
- **TEMPO CC (Coliform Count)**
- **Soleris NF-TVC**
- **Soleris Coliform Test**
- **Soleris Direct Yeast & Mold**
- **CompactDry "Nissui" YMR**
- **Quant X Fungal One Step**
- **DetectX Combined**

Confirmation Testing:

- **GENE-UP® EHEC Series**
- **BAX System Real-Time PCR Assay Suite for STEC**
- **iQ-Check Salmonella II Real-Time PCR**
- **iQ-Check STEC VirX/SerO/SerOII**
- **GENE-UP Salmonella 2 (SLM2)**
- **PathoSEEK Salmonella and STEC E.coli Multiplex Assay with SenSATIVax Extraction**
- **3M Molecular Detection Assay 2-Salmonella**
- **3M Molecular Detection Assay 2-STE C Gene Screen (stx and eae)**
- **3M Molecular Detection Assay 2-STE C Gene Screen (stx)**
- **GENE-UP PRO STEC/Salmonella Assay**
- **BAX System Real-Time PCR Assay for E.coli 0157:H7 Extract**

Sample Analysis (Water Activity):

- **Standard Practice for Determination of Water Activity in Cannabis Flower: ASTM D8196**

HEAVY METALS:

For method validations, please incorporate the Standard Method Performance Requirements (SMPRs) listed below:

- **Determination of Heavy Metals in a Variety of Cannabis and Cannabis Derived Products: AOAC SMPR 2020.001**

Sample Analysis:

- **Heavy Metals in a Variety of Cannabis and Cannabis Derived Products: AOAC 2021.03**

REFERENCES

- Freeman K, McHenry M, Cats-Baril W, Grace T. Cannabis Testing for Public Safety-Best Practices for Vermont Analytical Laboratories. PhytoScience Institute, 2016 January.
- Upton R, Craker L, ElSohly M, Romm A, Russo E, Sexton M. Cannabis Inflorescence: Standards of Identity, Analysis, and Quality Control. American Herbal Pharmacopoeia. Revision 2014.
- Thiex N, Ramsey C, Bhikha R, Cook J, Crawford A, Danielson D, Graves Q, Hirt W, Hoffman G, Andujar L, McLaughlin M, McManus K, McMullen S, Montalbano A, Novotny L, Salfinger Y, Schroder B, Scott T, Stephens R. Guidance on obtaining defensible samples. GOODSamples: Sampling and Sample Handling Working Group. 2015 October.
- Farrer DG. Technical Report: Oregon Health Authority's process to decide which types of contaminants to test for in cannabis. Oregon Health Authority. 2015 December.
- Osterbauer N, Krepps S, Sackett J, Holladay C, Wendt E, Wells D, Price S, Kristof J. Protocol for Collecting Samples of Usable Marijuana. ORELAP-SOP-001 Rev.3.0. 2016 December.
- Sexton M, Ziskind J. Sampling Cannabis for Analytical Purposes. BOTEC Analysis Corp. 2013 November.
- Voelker R, Holmes M. Pesticide use on cannabis. Cannabis Safety Institute, 2015 June.
- Holmes M, Vyas J, Steinbach W, McPartland J. Microbiological Safety Testing of Cannabis. Cannabis Safety Institute, 2015 May.
- International Council for Harmonisation. Guideline Q3C (R5) on Impurities: Guidelines for residual solvents. 2015 September.
- International Council for Harmonisation. Guideline Q3D Elemental Impurities: Guidance for Industry. 2015 September.
- Unger P, Brauinger R, Hudalla C, Holmes M, Sherman B. Standards for Cannabis Testing Laboratories. Cannabis Safety Institute. 2014 December.
- Raber J, Elzings S, Kaplan C. Understanding dabs: contamination concerns of cannabis concentrates and cannabinoid transfer during the act of dabbing. The Journal of Toxicological Sciences. Vol. 40, No. 6, 797-803, 2015 September.
- Aldous K, Applen J, Fan Z, Fox M, Kassner S, Latshaw M, Nascarella M, Starr G, Swantek S, Verbrugge D. Guidance for State Medical Cannabis Testing Programs. Association of Public Health Laboratories. 2016 May.
- International Council for Harmonisation. Guidance for Industry: Q1A(R2) Stability Testing of New Drug Substances and Products. 2003 November.
- Tchounwou P, Yedjou C, Patlolla A, Sutton D. Heavy Metals Toxicity and the Environment. Molecular, Clinical and Environmental Toxicology. Volume 101 of EXS, 133-164. 2012 April.
- U.S. Pharmacopeia Convention (USP). Chemical Tests / 561 Articles of Botanical Origin. 2014 July.
- Agricultural Improvement Act of 2018, Pub. L. No. 115 – 334, 132 Stat. 4490 (2018).
<https://www.congress.gov/115/plaws/publ334/PLAW-115publ334.pdf>